Adolescent Behaviors and Experiences Survey						
	Heterosexual		Gay, Lesbian, or Bisexual		Other/Questioning	
Alcohol and Other Drug Use	**Percentage**	**Confidence Interval**	**Percentage**	**Confidence Interval**	**Percentage**	**Confidence Interval**
Had their first drink of alcohol before age 13 years (other than a few sips)	9.5	8.2 - 11.1	19.4	15.7 - 23.7	14.8	11.2 - 19.4
Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	18.8	15.6 - 22.5	26.4	22.1 - 31.3	20.1	15.0 - 26.4
Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey)	7.8	5.6 - 10.8	9.0	6.5 - 12.5	7.5	4.8 - 11.7
Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	2.4	1.6 - 3.5	2.1	1.1 - 3.9	0.6	0.2 - 1.7
Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	37.3	32.8 - 42.0	44.4	36.1 - 53.0	36.8	25.8 - 49.4
Ever used marijuana (one or more times during their life)	23.5	21.1 - 26.0	35.7	31.4 - 40.3	25.8	22.0 - 29.9
Tried marijuana for the first time before age 13 years	2.9	2.2 - 3.9	6.1	4.4 - 8.4	4.4	2.6 - 7.2
Currently used marijuana (one or more times during the 30 days before the survey)	12.0	10.3 - 13.8	18.5	14.7 - 22.9	12.1	9.1 - 15.8
Ever used synthetic marijuana (one or more times during their life)	5.1	4.0 - 6.5	8.1	5.9 - 10.9	6.7	4.5 - 9.7
Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	8.0	6.9 - 9.2	16.0	13.1 - 19.4	15.5	12.2 - 19.5
Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)	3.8	3.0 - 4.7	6.8	5.2 - 9.0	5.4	3.8 - 7.8
Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	1.2	0.8 - 1.7	1.9	1.1 - 3.3	1.7	0.8 - 3.8
Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	4.0	3.2 - 5.0	13.7	10.6 - 17.5	10.1	6.9 - 14.5
Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.4	0.2 - 0.8	1.1	0.5 - 2.3	0.9	0.4 - 2.0
Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	0.5	0.3 - 0.9	1.5	0.8 - 2.9	1.4	0.5 - 3.4
Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)	1.8	1.2 - 2.6	1.9	1.0 - 3.7	3.1	1.5 - 6.1
Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)	3.1	2.3 - 4.3	6.0	4.2 - 8.6	5.8	3.9 - 8.6
Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times during their life)	7.4	6.1 - 9.0	17.5	14.3 - 21.3	12.5	9.0 - 17.2
Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	0.3	0.2 - 0.6	0.9	0.4 - 2.0	0.8	0.3 - 2.1
Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended school in-person during the 12 months before the survey)	8.5	6.4 - 11.3	11.3	8.2 - 15.2	14.4	10.4 - 19.5