

Adolescent Behaviors and Experiences Survey						
	Heterosexual		Gay, Lesbian, or Bisexual		Other/Questioning	
Mental Health	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval
Who felt sad or hopeless (almost every day for two or more weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	36.7	34.1 - 39.4	75.7	70.9 - 79.9	68.7	63.6 - 73.4
Seriously considered attempting suicide (during the 12 months before the survey)	13.6	11.7 - 15.8	46.8	41.5 - 52.2	39.5	34.6 - 44.7
Made a plan about how they would attempt suicide (during the 12 months before the survey)	9.9	8.3 - 11.8	37.9	34.3 - 41.8	32.3	27.4 - 37.5
Actually attempted suicide (one or more times during the 12 months before the survey)	5.2	4.2 - 6.5	26.3	21.8 - 31.4	16.5	11.8 - 22.7
Had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	0.9	0.7 - 1.3	5.9	3.7 - 9.4	3.6	2.1 - 6.0
Reported that their mental health was most of the time or always not good (including stress, anxiety, and depression, during the 30 days before the survey)	25.5	22.5 - 28.8	54.9	49.5 - 60.2	45.7	40.5 - 50.9