Adolescent Behaviors and Experiences Survey						
	Heterosexual		Gay, Lesbian, or Bisexual		Other/Questioning	
Sexual Behaviors	**Percentage**	**Confidence Interval**	**Percentage**	**Confidence Interval**	**Percentage**	**Confidence Interval**
Ever had sexual intercourse	28.6	26.0 - 31.3	33.4	28.4 - 38.9	23.9	19.6 - 28.9
Had sexual intercourse for the first time before age 13 years	1.6	1.2 - 2.2	5.1	3.5 - 7.2	2.6	1.5 - 4.8
Had sexual intercourse with four or more persons during their life	5.6	4.5 - 7.0	7.0	4.9 - 9.8	5.3	3.3 - 8.4
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	19.5	17.1 - 22.2	19.7	15.5 - 24.7	12.4	9.3 - 16.2
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	14.9	11.7 - 18.8	21.4	14.4 - 30.5	26.7	15.3 - 42.3
Used a condom during last sexual intercourse (among students who were currently sexually active)	58.3	54.6 - 61.9	35.7	26.7 - 45.9	41.4	31.4 - 52.2
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	21.6	18.3 - 25.4	23.3	16.8 - 31.3	29.3	18.0 - 43.8
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	5.0	4.4 - 5.8	10.1	6.7 - 14.8	6.6	4.6 - 9.5
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	4.3	3.5 - 5.1	7.8	5.0 - 11.9	3.5	1.9 - 6.3
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	5.8	4.2 - 7.9	7.8	3.8 - 15.4	5.0	1.6 - 14.6
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	33.5	29.7 - 37.6	35.2	26.7 - 44.8	39.6	27.6 - 53.1
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	12.5	10.4 - 14.9	8.2	4.5 - 14.7	11.2	4.9 - 23.3
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	12.8	10.7 - 15.3	18.8	11.1 - 30.1	9.4	5.2 - 16.4