



Click to learn about *Clostridioides difficile* infections

Topic

Number of Cases

Case Rates

Strain Type

Year

All

View By

Epi Class

Age

☒ Select all

☒ CA - 1-17 years

☒ CA - 18-44 years

☒ CA - 45-64 years

☒ CA - 65+ years

☒ HA - 1-17 years

☒ HA - 18-44 years

☒ HA - 45-64 years

☒ HA - 65+ years



Download Data

Candida bloodstream infections (Candidemia)

Clostridioides difficile infections

Invasive *Staphylococcus aureus* infections


Multi-site Gram-negative Surveillance Initiative

HAIC Viz



Clostridioides difficile infections in HAIC surveillance area

Epi Class by age



Note: Epidemiologic classification (epi class): A CDI case is classified as community-associated (CA) if the *Clostridioides difficile*-positive stool specimen was collected on an outpatient basis or within 3 days after hospital admission in a person with no documented overnight stay in a healthcare facility in the preceding 12 weeks. All CDI cases that do not meet the aforementioned criteria are classified as healthcare-associated (HA).

[Surveillance areas](#) have changed over time.



Click to learn about *Clostridioides difficile* infections

Topic

- Number of Cases
- Case Rates
- Strain Type

Year

All

View By

- All cases
- Total

- ☒ Select all
- ☒ Total



Download Data

Candida bloodstream infections (Candidemia)

Clostridioides difficile infections


Invasive *Staphylococcus aureus* infections

Multi-site Gram-negative Surveillance Initiative

HAIC Viz



Rates of *Clostridioides difficile* infections in HAIC surveillance area



Note: Epidemiologic classification (epi class): A CDI case is classified as community-associated (CA) if the *Clostridioides difficile*-positive stool specimen was collected on an outpatient basis or within 3 days after hospital admission in a person with no documented overnight stay in a healthcare facility in the preceding 12 weeks. All CDI cases that do not meet the aforementioned criteria are classified as healthcare-associated (HA). [Surveillance areas](#) have changed over time.



Click to learn about *Clostridioides difficile* infections

Topic

- Number of Cases
- Case Rates
- Strain Type

Year

All

View By

- Ribotype
- Select all
 - CA - 002
 - CA - 014
 - CA - 020
 - CA - 027
 - CA - 106
 - HA - 002
 - HA - 014
 - HA - 020
 - HA - 027
 - HA - 106

Download Data

Candida bloodstream infections (Candidemia)


Clostridioides difficile infections

Invasive Staphylococcus aureus infections

Multi-site Gram-negative Surveillance Initiative

HAIC Viz

Clostridioides difficile ribotypes by epidemiologic classification in HAIC surveillance area



Note: Epidemiologic classification (epi class): A CDI case is classified as community-associated (CA) if the *Clostridioides difficile*-positive stool specimen was collected on an outpatient basis or within 3 days after hospital admission in a person with no documented overnight stay in a healthcare facility in the preceding 12 weeks. All CDI cases that do not meet the aforementioned criteria are classified as healthcare-associated (HA).

[Surveillance areas](#) have changed over time.